

2020 P4P Best Practice Guidelines

**\*\*ENSURE THAT EVERY PATIENT VISIT IS DOCUMENTED AND BILLED\*\***

Measure	Who it applies to	Frequency	Qualified Event(s) and accepted codes	
BREAST CANCER SCREENING (BCS)	Females 50-74 yrs	Every 2 years	Mammogram	
CHLAMYDIA SCREENING IN WOMEN (CHL)	Females 16-24 yrs identified as sexually active	Annually	Chlamydia test (87110, 87270, 87320, 87490-87492, 87810)	
COLORECTAL CANCER SCREENING (COL)	Males and Females 50-75 yrs	Annually	FOBT (82270, 82274)	
		Every 5 years	Flexible Sigmoidoscopy	
		Every 10 years	Colonoscopy	
EVIDENCE-BASED CERVICAL CANCER SCREENING (ECS)	Females 21-65 yrs and 67+  (66 yr olds are excluded due to the 3 yr look back period. Depending on where their bday falls, they fall into Appropriately Screened and Screened too Frequently)	Every 3 years	1 Pap test: females 21-65 yrs w/ no hysterectomy (88141-88143, 88147, 88148, 88150, 88152-88155, 88164-88167, 88174, 88175)	
		Every 5 years	1 Pap test: females 21-65 yrs w/ no hysterectomy AND an HPV test tested at the same time	
		None	No Pap test: females 21-65 yrs with a hysterectomy and females 67+	
		Excluded	Females who have been DX with HPV or Cervical Cancer are excluded from this measure ICD-10 (Cervical Cancer) C53.0, C53.1, C53.8, C53.9 ICD-10 (HPV) R87.810, R87.811, R87.820, R87.821	
IMMUNIZATIONS FOR ADOLESCENTS (IMA)	Males and Females who turn 13 in 2015	1 between 11 and 13 yrs	Meningococcal (90733, 90734)	
		1 between 10 and 13 yrs	Tdap (90715) OR Td (90714, 90718)	
		3	HPV vaccinations between 9 and 13 (90649 or 90650)	
USE OF IMAGING STUDIES FOR LOW BACK PAIN (LBP)	Members 18-50 yrs with primary dx of low back pain		<b>DID NOT</b> have an imaging study (MRI, x-ray or CT scan) within 28 days of dx	
ANNUAL MONITORING FOR PATIENTS ON PERSISTENT MEDICATIONS (MPM)	Members 18+ yrs who rcvd at least a 180 treatment days of one of the following: ACE Inhibitors or ARBs, Digoxin or Diuretics	Annually	1 of the following: a lab panel test, a serum potassium and serum creatinine, or a serum potassium and blood urea nitrogen	
PROPORTION OF DAYS COVERED BY MEDICATIONS (PDC)	Males and Females 18+ who filled at least 2 prescriptions for one of the following:		Oral diabetes medication	Compliance is based on the proportion of days covered threshold of 80% for these medications
			Renin Angiotensin System Antagonists	
			Statin medications	
COORDINATED DIABETES CARE	Males and Females 18-75 yrs with diabetes (Type 1 and Type2)	Annually	Eye Exam	By eye care professional
			Monitoring Diabetic Nephropathy	Nephropathy Screening test, Urine microalbumin test, received medical attention for Nephropathy, OR ACE inhibitor/ARB therapy
			HbA1c Control	<8.0%
			Blood Pressure reading (result of <140/80)	Systolic: 3074F, 3075F, 3077F; Diastolic: 3078F, 3079F, 3080F
CHILDHOOD IMMUNIZATION STATUS (CIS)	Males and Females who turn 2 in 2015 and have received the following:	4	DTaP	90698, 90700, 90721, 90723
		3	IPV	90698, 90713, 90723
		1	MMR	90707 or 90710
		3	HiB	90645-90648-90698, 90721, 90748
		3	Hep B	90723, 90740, 90744, 90747, 90748
		1	VZV	90710 or 90716
		4	PCV	90669 or 90670
		1	Hep A	90633
2 or 3	Rotavirus	90681 or 90680		
Avoidance of Antibiotic Treatment for Adults w/ Acute Bronchitis (AAB)	Adults 18-64 yrs w/ dx of acute bronchitis		No antibiotic prescription given	
Asthma Medication Ratio (AMR)	Members 5-64 who were identified as having persistent asthma		Ratio of controller medications to total asthma medication of .50 or greater	Ratio = Units of Controller Meds divided by Units of Controller Meds + Units of Reliever Meds
Appropriate Testing for Children with Pharyngitis (CWP)	Children 2-18 who were diagnosed w/ pharyngitis		Dispensed an antibiotic and received a group A strep test	
Appropriate Treatment for Children with Upper Respiratory Infection (URI)	Children 3 mths-18 yrs given dx of upper respiratory infection		No antibiotic prescription given	